

**OBSERVATIONS OF THE Ca II IR TRIPLET IN HIGH LUMINOSITY
QUASARS: EXPANDING THE SAMPLE**

**M. L. Martínez-Aldama¹, D. Dultzin¹, P. Marziani², J. W. Sulentic³,
A. Bressan⁴ and G. M. Stirpe⁵**

¹*Instituto de Astronomía, Universidad Nacional Autónoma de México, Mexico*

²*INAF, Osservatorio Astronomico di Padova, Padova, Italia*

³*Instituto de Astrofísica de Andalucía (CSIC), Granada, España*

⁴*Astrophysics Sector, SISSA/ISAS, Trieste, Italia*

⁵*INAF, Osservatorio Astronomico di Bologna, Bologna, Italia*

E-mail: maldama@astro.unam.mx

We present spectroscopic observations of 10 quasars at intermediate redshift obtained with the Infrared Spectrometer and Array Camera (ISAAC) on the ESO Very Large Telescope, covering the region of OI 8446 and the CaII triplet 8498, 8542, 8662. The new observations almost double the size of the sample described in Martínez-Aldama et al. (2015) that contained almost all CaII observations previously available for intermediate z quasar. The new spectra cover CaII in one of the relatively rare, extreme Population A quasars, and allow for a more refined and statistically significant discussion on low-ionization line formation and recent episodes of star formation in intermediate redshift quasars.