

PROGRESS REPORT ON THE SAO OPTICAL MONITORING OF TYPE 1 AGN

**A. I. Shapovalova¹, L. Č. Popović^{2,3}, D. Ilić³, A. Kovačević³,
A. N. Burenkov¹ and V. H. Chavushian⁴**

¹*Special Astrophysical Observatory of the Russian Academy of Science, Nizhnij Arkhyz,
Karachaevo-Cherkesia 369167, Russia*

²*Astronomical Observatory, Volgina 7, 11060 Belgrade, Serbia*

³*Department of Astronomy, Faculty of Mathematics, University of Belgrade,
Studentski trg 16, 11000 Belgrade, Serbia*

⁴*Instituto Nacional de Astrofísica, Óptica y Electrónica, Apartado Postal 51,
CP 72000, Puebla, Pue. Mexico*

E-mail: dilic@math.rs

Here we give a progress report on the latest result of the long-term optical monitoring of type 1 active galactic nuclei (AGN). A campaign, which is coordinated by the Special Astrophysical Observatory (SAO) is performed using telescopes of SAO (Russia), INAOE and OAN-SPM (Mexico), with the main aim to use the broad line shape variability to constrain the physics and kinematics of the line emitting regions. In total a dozen of type 1 AGN are observed for decades, and here we outline the latest results, especially in case of a quasar E1821+643 and Seyfert 1 galaxy NGC 7469.