

**CHEMI-IONIZATION/RECOMBINATION PROCESSES  
IN THE BROAD-LINE REGION OF AGNs**

**V. A. Srećković<sup>1</sup>, M. S. Dimitrijević<sup>2,3</sup> and Lj. M. Ignjatović<sup>1</sup>**

<sup>1</sup>*Institute of Physics, Pregrevica 118, 11080 Zemun, Belgrade, Serbia*

<sup>2</sup>*Astronomical Observatory, Volgina 7, 11060 Belgrade, Serbia*

<sup>3</sup>*Sorbonne Université, Observatoire de Paris, Université PSL, CNRS, LERMA,  
F-92190 Meudon, France*

*E-mail: vlada@ipb.ac.rs, mdimitrijevic@aob.rs, ljuba@ipb.ac.rs*

The importance of Chemi-ionization processes in  $H^*(n) + H(1s)$  collisions in AGN BLR clouds has been investigated. These processes must have influence on ionization and the populations of hydrogen highly excited atoms in moderately ionized layers of dense parts of the BLR clouds. From the results it follows that these investigated processes are of interest for the research and modelling of such medium. The obtained results could be also useful for modeling of different stellar atmospheres, as well as for the investigation of Rydberg states of hydrogen and for the study of their influence during the cosmological recombination epoch (see e.g. Mihajlov et al. 2011).

**References**

Mihajlov, A. A., Ignjatović, L. M., Srećković, V. A., Dimitrijević, M. S.: 2011, *ApJS*, **2**, 193.